

What do we know so far of cosmic neutrinos?
Highlights of the cosmic discoveries


- Diffuse Cosmic Neutrino Flux
- Cosmic Sources
- Galactic Neutrinos
- The Unexpected

A neutrino beam 'for free'


Cosmic rays hit the atmosphere
=> particle showers

Particles decays
=> Muon/Electron neutrinos
created in ratio 2:1

Backgrounds for cosmic neutrino
searches, but useful
for neutrino oscillation studies



Flavors and neutrino/antineutrinos



Honda et al, Phys. Rev. D92 (2015) 023004

Conventional atmospheric neutrino flux

from kaon and pion decays

- Mainly muon neutrinos
- More neutrinos than anti-neutrinos
- Fluxes depend on atmospheric seasonal changes
-> slight annual flux changes

In addition 'prompt flux':

Neutrinos from decay of heavy mesons (D-mesons)

- Higher energies, harder energy spectrum
- First hint of this flux component seen by now in IceCube



Atmospheric background & cosmic neutrinos

~1km³ detector:

Atmospheric μ 10^9 per year

Atmospheric ν 10^4 per year

Cosmic ν ~100 per year




High energies -> Cosmic neutrinos dominate

In order to identify sources: Time / space coincidence with other cosmic probes


Diffuse cosmic neutrino flux

The two first cosmic neutrinos detected by IceCube




High energy events could not be explained by atmospheric background

One of the first extraterrestrial neutrino candidates
in IceCube - moved to Paris





Picture by Doug Cowen, Rencontres de Blois 2013

Active veto



Veto allows to cleanly identify
neutrino interactions





- Veto allows to cleanly identify neutrino interactions
- Reduces atmospheric neutrino 'background' from above due to signals of high-energy muon accompanying neutrino production

Gaisser et al, PhysRevD, 90:023009 (2014)

Led to discovery of many more cosmic neutrino candidates
HESE: High Energy Starting Events



High energy diffuse neutrino flux

Many more searches established the cosmic diffuse neutrino flux




High energy diffuse neutrino flux

Many more searches established the cosmic diffuse neutrino flux




	$-2\Delta\ln\mathcal{L}$ compared to SPL
Single PL	0
SPL + Cutoff	1.8 (0.9 σ)
Log Parabola	19.8 (4.2 σ)
SPL + Bump	22.3 (3.9 σ)
Broken PL	27.3 (4.7 σ)

New: A 'broken power law' clearly identified



Z. Chen, Z. Zhang, J. Kiryluk (for IceCube)
PoS(ICRC2025)1011



GVD, arXiv.2507.01893

Potential differences could be expected from different event samples:

- Different sky regions probed
- Different energy ranges probed
- Different flavors

Different source scenarios

$$p + \gamma \rightarrow \Delta^+ \begin{cases} \rightarrow p + \pi^0 \\ \rightarrow n + \pi^+ \end{cases}$$

$$\pi^0 \rightarrow \gamma\gamma$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu$$

$$\quad \quad \quad \hookrightarrow e^+ + \bar{\nu}_\mu + \nu_e$$

$$n \rightarrow e^- + p + \bar{\nu}_e \quad \text{Neutron lifetime: 881 s}$$

$$p + p \rightarrow \pi^\pm + X$$

$$\quad \quad \quad \hookrightarrow \mu^\pm + \nu_\mu (\bar{\nu}_\mu)$$

$$\quad \quad \quad \hookrightarrow e^\pm + \bar{\nu}_\mu (\nu_\mu) + \nu_e (\bar{\nu}_e)$$

Different cosmic source scenarios

$$\nu_e : \nu_\mu : \nu_\tau$$

'Standard'

-> charged pion decay, muon decay

$$1 : 2 : 0$$

Muon damped source

-> strong magnetic fields, muon decay suppressed

-> pion decay dominant

$$0 : 1 : 0$$

Neutron beam source

-> extremely strong magnetic field

-> cosmic rays heavy nuclei



$$1 : 0 : 0$$

Note: Also different neutrino/antineutrino ratios

Neutrino flavor ratio at Earth



- Different astrophysical scenarios will lead to different observed flavor compositions
- Potential to find new physics effects

- Flavor ratio can be energy dependent
- In neutron decay scenario only antineutrinos
neutrino telescopes don't distinguish neutrinos and antineutrinos, but Glashow resonance occurs only for antineutrinos



Neutrino flavor ratio at Earth


- Different astrophysical scenarios will lead to different observed flavor compositions
- Potential to find new physics effects




Neutrino flavor ratio at Earth

'Standard' phase space


-> Deviations indicate new physics




IceCube measurement




Prospects for measurements – energy dependence



Cosmic neutrino sources




22 September 2017
Blazar TXS 0506 in flaring state
-> High energy neutrino detected




Optical sky observations around the
IceCube observation

Blazar TXS 0506+056

Angular resolution crucial to identify
neutrino sources



γ ray observations



Puzzling: Neutrino coincidence with γ ray alert,
but no enhanced γ ray flaring in 2014

γ ray absorption?

3 November 2022 Announcement by IceCube,

Publication in 'Science' journal


Evidence ($>4\sigma$) of neutrino emission from NGC 1068 (Messier 77)



IceCube events around NGC 1068C

Events close to NGC 1068

Can you spot a source with
80 events (excess over background)?



Evidence for neutrino emission from the nearby active galaxy NGC1068, *SCIENCE* 2022, Vol 378, Issue 6619


Dataset. DOI: [10.21234/03fq-rh11](https://doi.org/10.21234/03fq-rh11)

IceCube events around NGC 1068C

Events close to NGC 1068

Can you spot a source with
80 events (excess over background)?

Red: Events with $\log(E/\text{GeV}) > 3.5$



Evidence for neutrino emission from the nearby active galaxy NGC1068, *SCIENCE* 2022, Vol 378, Issue 6619


Dataset. DOI: [10.21234/03fq-rh11](https://doi.org/10.21234/03fq-rh11)

IceCube events around NGC 1068C

Events close to NGC 1068

Can you spot a source with
80 events (excess over background)?


Red: Events with $\log(E/\text{GeV}) > 3.5$



Evidence for neutrino emission from the nearby active galaxy NGC1068, *SCIENCE* 2022, Vol 378, Issue 6619

Dataset. DOI: [10.21234/03fq-rh11](https://doi.org/10.21234/03fq-rh11)


IceCube events around NGC 1068



Evidence for neutrino emission from the nearby active galaxy NGC1068, *SCIENCE* 2022, Vol 378, Issue 6619

Dataset. DOI: [10.21234/03fq-rh11](https://doi.org/10.21234/03fq-rh11)

IceCube events around NGC 1068



Evidence for neutrino emission from the nearby active galaxy NGC1068, *SCIENCE* 2022, Vol 378, Issue 6619
Dataset. DOI: [10.21234/03fq-rh11](https://doi.org/10.21234/03fq-rh11)

Search for event excess by use of likelihood evaluation:

- Maximize likelihood L at each point in the sky
- Signal and background PDFs contain spatial and energy terms


$$L(n_s, x_s, \gamma) = \prod_i^{events} \left(\frac{n_s}{N} S_i(|x_i - x_s| \sigma_i, E_i, \gamma) + \frac{N - n_s}{N} B_i(\delta_i, E_i) \right)$$

Spatial term for likelihood was:


$$S_i(|\vec{x}_i - \vec{x}_s|, \sigma_i) = \frac{1}{2\pi\sigma_i^2} \exp\left(-\frac{|\vec{x}_i - \vec{x}_s|^2}{2\sigma_i^2}\right)$$

Assumes here Gaussian shape of resolution,
but can be replaced also by non-gaussian shape

Refined resolution for the NGC 1068 analysis



IceCube events around NGC 1068





1) Evidence for neutrino emission from the nearby active galaxy NGC1068, *SCIENCE* 2022, Vol 378, Issue 6619
Dataset. DOI: [10.21234/03fq-rh11](https://doi.org/10.21234/03fq-rh11)

[2] IceCube Search for Neutrino Emission from X-ray Bright Seyfert Galaxies, 2025 *ApJ* **988** 141

[3] Search for neutrino emission from hard X-ray AGN with IceCube, 2025 *ApJ* **981** 131

[4] Evidence for Neutrino Emission from X-ray Bright Active Galactic Nuclei with IceCube, *arXiv:2510.13403*



Correlation with X-ray flux/luminosity



Galactic neutrino flux


One of the first neutrino skies from IceCube

7.5 years data (102 events)




$$TS = -2\Delta \ln(\mathcal{L})$$

2017 paper




2019 paper




2 years of data
Galactic Plane p-value: 65%

7 years of data
Galactic Plane p-value: 2.1% (2σ)



- Deep learning improved resolution by 2, sensitivity by 3
- 13 years of data Galactic Plane: 5.7σ post trial (Lepton-Photon 2025, Madison)




<http://adc.gsfc.nasa.gov/mw>

Milky Way
viewed in
neutrinos



Halzen,
Neutel 2025

Discovery horizon for $L_{100\text{TeV}} = 10^{34} \text{ erg/s}$ ($\Phi \propto E^{-2}$)



- Complementary galactic regions to be probed by future detectors

The Unexpected

The international journal of science / 13 February 2025

nature

COSMIC CATCHER



Deep-sea telescope detects
neutrino with highest energy
ever recorded

Nikhef





Universiteit
Leiden

Vol. 638, No. 8020
£10.00 nature.com




Unique event



A moving detector

Tilt Compared to tilt of tower of Pisa?

Tilt amplitude



Tower of Pisa:
~4 degrees



4.5deg

1 year of data

- Detector positions monitored every 10 minutes with acoustic signals
- Position accuracy: ~20cm

A very well reconstructed muon


Intrinsic reconstruction uncertainty: $\sim 0.1^\circ$



IceCube


High energy cascade 'Bert'

- Likelihood fit on the waveforms
- Shaded areas disregarded (saturation/ systematics)
- High energy events direction resolution 7-25 degrees



Unique Event


- Light profile consistent with at least 3 large energy depositions along the muon track
- Characteristic of stochastic losses from very high energy muons



Unique Event

Simulation

Aantal PMTs




↑ direction – $\cos \theta$ ↓

Red: atmospheric muons


Blue: cosmic neutrinos

Data




↑ direction – $\cos \theta$ ↓

Energy estimate



- Energy loss simulated for **10 PeV** and **100 PeV** muons with the very same location
- 3672 PMTs (35%) were triggered in the detector
- Muons simulated at 10 PeV almost never generate this much light

Energy estimate



- Estimate of the muon energy:



$$E_{\mu} = 120^{+110}_{-60} \text{ PeV}$$

- Estimate of neutrino energy:

$$E_{\nu} = 220^{+570}_{-100} \text{ PeV}$$



(assumption: E^{-2} spectrum)

Energy estimate




Where did it come from?

Horizontal muon from a neutrino interaction




It can't be an atmospheric muon


Distance travelled by 100 EeV muons



Less than 40km!



Simulations: Isotropic 100 PeV neutrino flux



Most likely detection
Is from a horizontal direction!



Direction – $\cos \theta$






**KM3NeT
Sites**

2.5km depth

3.5km depth




International Bathymetric Chart of the Mediterranean

<https://www.ngdc.noaa.gov/mgg/ibcm/ibcmbath.html>


How accurate do we know the absolute orientation/tilt of the detector?

3.5 km deep!




- Transfer of GPS coordinates from Sea Surface during deployments with acoustic signals
- Comparison of 2 existing bathymetric data sets
- Check average tilt of the lines
- Evaluation of the cosmic ray shadow of the moon/sun

Check the shadow of the moon/sun in cosmic rays






Probing independent rotations




Credit: Jelle Oonk, Leiden University




Effects of Misalignment





Effects of Misalignment




Check the shadow of the moon/sun in cosmic rays




Confirmation of good data filtering, calibration, reconstruction!



Pointing on the sky - suspect sources (1): Blazars



Pointing on the sky - suspect sources (1): Blazars



- Large uncertainty region
- Intrinsic reconstruction uncertainty $\sim 0.1^\circ$
- Current uncertainty dominated by uncertainty in absolute detector orientation



- New acoustic beacon deployments with high accuracy (1m) will enable to measure absolute orientation
- Moon shadow analysis will serve as cross check

=> **We will in the future improve on the direction information of this event!**

KM3-230213A Blazar candidates

#6 0605-085


Fermi-LAT
γ-ray observations
1-100 GeV



(a) The *Fermi*-LAT light curve and a VLBI image of 0605-085: the brightest radio source in the neutrino localization region that experiences a gamma-ray flaring activity around the neutrino arrival (Section 5.1).

#8: PMN J0606-0724


OVRO/RATAN
radio observations
15GHz



(b) The radio light curve for PMN J0606-0724 that experiences a major flare in close coincidence to the neutrino arrival (Section 5.2).


#1: MRC 0614-083

eROSITA/Swift-XRT
X-ray observations
0.2-2.3 keV



(c) The X-ray light curve for MRC 0614-083 that indicates a flaring activity around the neutrino arrival (Section 5.3).

Pointing on the sky - suspect sources (2): Galactic?



No obvious suspect, generally hard to achieve this high energy


Unexplored energy regime

LHC@CERN

Previous highest
energy neutrino

This observation

Intensity of neutrinos




Neutrino energy (logarithmic scale)


LHC@CERN

Previous highest
energy neutrino

This observation




Unexplored energy regime



Best-fit flux taking into account non-observations from IceCube and Auger

Varying the cosmogenic neutrino models



Moving forward

Many neutrino observatories substantially growing, coming online

- Angular resolution crucial
 - > expect improvements from refined reconstructions & lower intrinsic uncertainties (water)
- Multi - messenger observations key to unravel the actual high energy cosmic sources
 - > connection between observatories essential

The neutrino sky has surprised us several times before
-> exciting decades ahead!



*'Draw me a neutrino' contest 2020
winner:: Evangelos Zacharopoulos*